arXiv:1608.06088v2 [math.DG] 9 Sep 2016

TRIHARMONIC RIEMANNIAN SUBMERSIONS FROM 3-DIMENSIONAL MANIFOLDS OF CONSTANT CURVATURE

TOMOYA MIURA AND SHUN MAETA

Abstract. For biharmonic maps, there is a famous conjecture named Chen's conjecture. In later paper, Wang and Ou gave an affirmative partial answer to submersion version of Chen's conjecture. In this paper, we give an affirmative partial answer to submersion version of generalized Chen's conjecture, that is, triharmonic Riemannian submersions from a 3-dimensional space form into a surface is harmonic.

1. Introduction

A harmonic map which is a critical point of the energy functional $E(\varphi) = \frac{1}{2}\int_M |d\varphi|^2 v_g$ is important notion in geometry.

In 1964, Eells and Sampson [9] extended a harmonic map to a polyharmonic map of order k. A polyharmonic map of order k is defined a critical point of the k-energy functional (cf. [8], [9])

$$E_k(\varphi) = \frac{1}{2}\int_M |(d+\delta)^k \varphi|^2 v_g. \tag{1.1}$$

For $k = 2$, the map called a biharmonic map and for $k = 3$, the map called a triharmonic map.

A fundamental study of biharmonic maps is to classify all biharmonic maps between Riemannian manifolds. For a biharmonic map, there is a famous conjecture.

Chen's conjecture (cf. [5]): *Any biharmonic isometric immersion $\varphi : (M^m, g) \to \mathbb{R}^n$ into the Euclidean space is harmonic.*

There are several affirmative partial answers to this conjecture. In particular, for M^2 in $\mathbb{E}^3$ [5] and for M^3 in $\mathbb{E}^4$ [6], [11], this conjecture

2010 *Mathematics Subject Classification.* primary 58E20, secondary 53C43.

Key words and phrases. harmonic maps, biharmonic maps, triharmonic maps, Riemannian submersions.

The second author is partially supported by the Grant-in-Aid for Young Scientists(B), No.15K17542, Japan Society for the Promotion of Science.

was proved. Also, for φ is a curve [7] and φ is proper [1], this conjecture was proved (see also [15], [13]).

For the Chen's conjecture, the second author posed the generalized Chen's conjecture.

Maeta's Generalized Chen's conjecture (cf. [14], see also [4]): *The only k-harmonic submanifolds (polyharmonic submanifolds of order k) in Euclidean spaces are the minimal ones.*

In later paper, Wang and Ou [22] showed the following.

Theorem 1.1 ([22])**.** *Let $\varphi : (M^3(c), g) \to (N^2, h)$ be a biharmonic Riemannian submersion from a space form of constant sectional curvature c. Then, φ is harmonic.*

We pose submersion version of generalized Chen's conjecture.

Conjecture 1.2.

1. *Are biharmonic submersions $\varphi : M^m(c) \to N^{m-1}$ harmonic?*
2. *Are triharmonic submersions $\varphi : M^m(c) \to N^{m-1}$ harmonic?*

In this paper, we give an affirmative partial answer to submersion version of generalized Chen's conjecture.

Theorem 1.3. *Let $\varphi : (M^3(c), g) \to (N^2, h)$ be a triharmonic Riemannian submersion from a space form of constant sectional curvature c. Then, φ is harmonic.*

Remark 1.4. *When we extend biharmonic immersions to triharmonic immersions, we have to add some additional assumptions (see* [16]*). However, in the above theorem, we do not need any additional assumptions.*

From the definition of a harmonic morphism (cf. [3]), we have the following.

Corollary 1.5. *Let $\varphi : (M^3(c), g) \to (N^2, h)$ be a triharmonic Riemannian submersion from a space form of constant sectional curvature c. Then, φ is harmonic morphism.*

By the same argument as in [22], we obtain the following result.

Corollary 1.6. (1) *If $\varphi : \mathbb{R}^3 \to (N^2, h)$ is a triharmonic horizontally homothetic submersion from Euclidean space, then (N^2, h) is flat and φ is a composition of an orthogonal projection $\mathbb{R}^3 \to \mathbb{R}^2$ followed by a covering map $\mathbb{R}^2 \to (N^2, h)$;*
(2) *There exists no triharmonic Riemannian submersion $\varphi : \mathbb{H}^3 \to (N^2, h)$ no matter what (N^2, h) is.*

2. Preliminaries

In this section, to proof the main theorem, we prepare several definitions and notions. In this paper, all manifolds, maps and tensor fields are assumed to be smooth. ∇ and ∇^N denote the Levi-Civita connections on (M, g) and (N, h), respectively. ∇^φ denotes the induced connection on $\varphi^{-1}TN$. The first variation formula of the trienergy is given as follows:

$$\tfrac{d}{dt}\Big|_{t=0} E_3(\varphi_t) = -\int_M \langle \tau_3(\varphi), V\rangle v_g, \quad (V \in \Gamma(\varphi^{-1}TN)),$$

where,

$$\begin{aligned}
\tau_3(\varphi) &= J(\overline{\triangle}(\tau(\varphi))) - \sum_{i=1}^m R^N(\nabla^\varphi_{e_i}\tau(\varphi), \tau(\varphi))d\varphi(e_i),\\
J(\tau(\varphi)) &= \overline{\triangle}(\tau(\varphi)) - \sum_{i=1}^m R^N(\tau(\varphi), d\varphi(e_i))d\varphi(e_i),\\
\overline{\triangle}\tau(\varphi) &= \sum_{i=1}^m \nabla^\varphi_{e_i}\nabla^\varphi_{e_i}\tau(\varphi) - \nabla^\varphi_{\nabla_{e_i}e_i}\tau(\varphi),
\end{aligned}$$

and

$$R^N(U, W) = \nabla^N_U\nabla^N_W - \nabla^N_W\nabla^N_U - \nabla^N_{[U,W]}. \quad (U, W \in \Gamma(TN)).$$

$\tau^3(\varphi)$ is called the *tritension field* of φ. A smooth map φ of (M^m, g) to (N^n, h) is said to be *triharmonic* if $\tau^3(\varphi) = 0$.

We introduce main tools of Riemannian submersions from 3-manifolds named adapted to a Riemannian submersion and integrability data (cf. [22]). Let $\varphi : (M^3, g) \to (N^2, h)$ be a Riemannian submersion. We can choose the special orthonormal frame adapted to φ $\{e_1, e_2, e_3\}$ on (M^3, g), where e_3 is vertical. The frame called adapted to Riemannian submersion. As is well known, the frame always exists. Since e_3 is vertical, $[e_1, e_3]$ and $[e_2, e_3]$ are vertical. Let $\{\epsilon_1, \epsilon_2\}$ be the orthonormal frame on (N^2, h). If we assume that

$$[\epsilon_1, \epsilon_2] = F_1\epsilon_1 + F_2\epsilon_2, \quad (F_1, F_2 \in C^\infty(N)),$$

then, we can assume

$$\begin{cases} [e_1, e_3] = k_1e_3, \\ [e_2, e_3] = k_2e_3, \\ [e_1, e_2] = f_1e_1 + f_2e_2 - 2\sigma e_3, \end{cases} \tag{2.1}$$

where, $k_1, k_2, \sigma, f_1 = F_1 \circ \varphi$ and $f_2 = F_2 \circ \varphi \in C^\infty(M)$ called integrability data of the adapted to Riemannian submersion φ.

3. Proof of the main theorem

Proposition 3.1. *Let $\varphi : (M^3, g) \to (N^2, h)$ be a Riemannian submersion with the adapted frame $\{e_1, e_2, e_3\}$ and the integrability data f_1, f_2, k_1, k_2 and σ. Then, the Riemannian submersion φ is triharmonic if and only if*

$$\begin{cases} \triangle^M A + f_1 e_1(B) + e_1(Bf_1) + f_2 e_2(B) + e_2(Bf_2) - f_1 k_1 B - k_2 f_2 B \\ -A\{K^N + f_1^2 + f_2^2\} + K^N(-e_2(k_1)k_2 + e_2(k_2)k_1 - k_1^2 f_2 - k_2^2 f_2) = 0, \\ \triangle^M B - f_1 e_1 A - e_1(Af_1) - f_2 e_2(A) - e_2(Af_2) + f_1 k_1 A + f_2 k_2 A \\ -B\{K^N + f_1^2 + f_2^2\} + K^N(e_1(k_1)k_2 - e_1(k_2)k_1 + f_1 k_2^2 + f_1 k_1^2) = 0, \end{cases} \tag{3.1}$$

where,

$$A = -\triangle^M k_1 - f_1 e_1(k_2) - e_1(k_2 f_1) - f_2 e_2(k_2) - e_2(k_2 f_2) + k_1 k_2 f_1 + f_2 k_2^2 + k_1 f_1^2 + k_1 f_2^2,$$

$$B = -\triangle^M k_2 + f_1 e_1(k_1) + e_1(k_1 f_1) + f_2 e_2(k_1) + e_2(k_1 f_2) - k_1 k_2 f_2 - f_1 k_1^2 + k_2 f_1^2 + k_2 f_2^2,$$

$$\triangle^M(k_1) = \sum_{i=1}^{3} e_i e_i(k_1) + f_1 e_2(k_1) - f_2 e_1(k_1) - k_1 e_1(k_1) - k_2 e_2(k_1),$$

and

$$K^N = R^N_{1212} \circ \pi = e_1(f_2) - e_2(f_1) - f_1^2 - f_2^2.$$

Proof A simple computation, using (2.1) and Koszul formula, yields

$$\begin{aligned} &\nabla_{e_1} e_1 = -f_1 e_2, \nabla_{e_1} e_2 = f_1 e_1 - \sigma e_3, \nabla_{e_1} e_3 = \sigma e_2, \\ &\nabla_{e_2} e_1 = -f_2 e_2 + \sigma e_3, \nabla_{e_2} e_2 = f_2 e_1, \nabla_{e_2} e_3 = -\sigma e_1, \\ &\nabla_{e_3} e_1 = -k_1 e_3 + \sigma e_2, \nabla_{e_3} e_2 = -\sigma e_1 - k_2 e_3, \nabla_{e_3} e_3 = k_1 e_1 + k_2 e_2. \end{aligned}$$

Furthermore, noting that e_3 is vertical, the following holds.

$$\begin{aligned} &\nabla^\varphi_{e_1} \epsilon_1 = -f_1 \epsilon_2, \quad \nabla^\varphi_{e_1} \epsilon_2 = f_1 \epsilon_1, \\ &\nabla^\varphi_{e_2} \epsilon_1 = -f_2 \epsilon_2, \quad \nabla^\varphi_{e_2} \epsilon_2 = f_2 \epsilon_1, \\ &\nabla^\varphi_{e_3} \epsilon_j = 0 \quad (j = 1, 2). \end{aligned}$$

Consequently, the tension field of φ is given by

$$\tau(\varphi) = -k_1 \epsilon_1 - k_2 \epsilon_2.$$

Under these preparations, we calculate $\tau^3(\varphi) = 0$. First, by a direct computation, we get

$$\overline{\triangle}\tau(\varphi) = A\epsilon_1 + B\epsilon_2.$$

Second, a straightforward computation yields

$$\begin{aligned} J(\overline{\triangle}\tau(\varphi)) = &[\triangle^M A + f_1 e_1(B) + e_1(Bf_1) + f_2 e_2(B) + e_2(Bf_2) \\ &- f_1 k_1 B - k_2 f_2 B - A(f_1^2 + f_2^2) - AK^N]\epsilon_1 \\ &+ [\triangle^M B - f_1 e_1(A) - e_1(Af_1) - f_2 e_2(A) - e_2(Af_2) \\ &+ f_1 k_1 A + f_2 k_2 A - B(f_1^2 + f_2^2) - BK^N]\epsilon_2. \end{aligned}$$

Third, in the same way, we get

$$\begin{aligned} -R^N(\nabla^\varphi_{e_i}\tau(\varphi), \tau(\varphi))d\varphi(e_i) = &K^N(-e_2(k_1)k_2 + e_2(k_2)k_1 - k_1^2 f_2 - k_2^2 f_2)\epsilon_1 \\ &+ K^N(e_1(k_1)k_2 - e_1(k_2)k_1 + k_1^2 f_1 + k_2^2 f_1)\epsilon_2. \end{aligned}$$

Therefore, we can get (3.1). □

If M^3 has a constant sectional curvature c, for all orthonormal frame $\{e_1, e_2, e_3\}$ on M^3 adapted to Riemannian submersion and for all integrability date f_1, f_2, k_1, k_2 and σ, we have two facts. (cf. [22]):

$$e_3(f_1) = e_3(f_2) = e_3(k_1) = e_3(k_2) = e_3(\sigma) = 0, \tag{3.2}$$

and we can choose adapted orthonormal frame $\{e_1, e_2, e_3\}$ such that $k_2 = 0$.

These facts simplify the triharmonic equation.

Corollary 3.2. *Let $\varphi : (M^3(c), g) \to (N^2, h)$ be a Riemannian submersion from a space form of constant sectional curvature c with an adapted frame $\{e_1, e_2, e_3\}$ and the integrability date f_1, f_2, k_1, k_2 and σ with $k_2 = 0$. Then, the Riemannian submersion φ is triharmonic if and only if*

$$\begin{cases} \triangle^M A' + f_1 e_1(B') + e_1(B'f_1) + f_2 e_2(B') + e_2(B'f_2) - f_1 k_1 B' \\ -A'\{K^N + f_1^2 + f_2^2\} - K^N(k_1^2 f_2) = 0, \\ \\ \triangle^M B' - f_1 e_1 A' - e_1(A'f_1) - f_2 e_2(A') - e_2(A'f_2) + f_1 k_1 A' \\ -B'\{K^N + f_1^2 + f_2^2\} + K^N(k_1^2 f_1) = 0, \end{cases} \tag{3.3}$$

where,
$A' = -\triangle^M k_1 + k_1 f_1^2 + k_1 f_2^2$,
and
$B' = f_1 e_1(k_1) + e_1(k_1 f_1) + f_2 e_2(k_1) + e_2(k_1 f_2) - f_1 k_1^2$.

By using Corollary 3.2, we prove the main theorem.

Theorem 3.3. *(Theorem 1.2.) Let $\varphi : (M^3(c), g) \to (N^2, h)$ be a triharmonic Riemannian submersion from a space form of constant sectional curvature c. Then, φ is harmonic.*

Proof By the facts, we choose an orthonormal frame $\{e_1, e_2, e_3\}$ adapted to the Riemannian submersion with integrability data f_1, f_2, k_1, k_2 and σ such that $k_2 = 0$. Since $M^3(c)$ has constant sectional curvature c, the following holds.

$$\begin{cases} R^M_{1312} = e_1(\sigma) - 2k_1\sigma = 0. \\ R^M_{1313} = -[-e_1(k_1) - \sigma^2 + k_1^2] = c. \\ R^M_{1323} = k_1 f_1 = 0. \\ R^M_{1212} = -(e_2(f_1) - e_1(f_2) + f_1^2 + f_2^2 + 3\sigma^2) = c. \\ R^M_{1223} = e_2(\sigma) = 0. \\ R^M_{2313} = e_2(k_1) = 0. \\ R^M_{2323} = -(-\sigma^2 + k_1 f_2) = c. \end{cases} \tag{3.4}$$

Here, by the 3rd equation in (3.4), if $k_1 = 0$ on $M^3(c)$, φ is harmonic. Therefore, we consider $k_1 \neq 0$ on some set $\Omega \subset M^3(c)$ (in this case, $f_1 = 0$ on Ω) and prove that the case cannot happen.

By the 1st equation of (3.4), we get

$$e_1(\sigma) = 2k_1\sigma. \tag{3.5}$$

By the 2nd and the 7th equations in (3.4), we get

$$e_1(k_1) = k_1^2 - k_1 f_2. \tag{3.6}$$

By applying e_1 to both sides of the 7th equation in (3.4), we get

$$e_1(f_2) = f_2^2 - k_1 f_2 + 4\sigma^2. \tag{3.7}$$

Also, in this case, we get

$$A' = -k_1^3 + 2f_2 k_1^2 + 4\sigma^2 k_1. \tag{3.8}$$

By using (3.5), (3.6), (3.7) and (3.8), the triharmonic equation is given as

$$\triangle^M A' - A'(e_1(f_2)) - (e_1(f_2) - f_2^2)(f_2 k_1^2) = 0. \tag{3.9}$$

Here, by (3.4) and (3.2), we get

$$e_2(A') = e_3(A') = 0.$$

Consequently, laplacian of A' can be rewritten as

$$\triangle^M(A') = e_1 e_1(A') - f_2 e_1(A') - k_1 e_1(A').$$

Since

$$e_1(A') = -3k_1^4 + 5f_2 k_1^3 + (-2f_2^2 + 28\sigma^2)k_1^2 - 4\sigma^2 f_2 k_1,$$

and

$$e_1 e_1(A') = -12k_1^5 + 22f_2 k_1^4 + (-10f_2^2 + 188\sigma^2)k_1^3 - 88\sigma^2 f_2 k_1^2 - 16\sigma^4 k_1,$$

the equation of (3.9) can be written as

$$-9k_1^5 + 19f_2k_1^4 + (164\sigma^2 - 9f_2^2)k_1^3 - 120\sigma^2 f_2 k_1^2 - 32\sigma^4 k = 0. \tag{3.10}$$

Since $k_1 \neq 0$ on Ω, (3.10) implies

$$-9k_1^4 + 19f_2k_1^3 + (164\sigma^2 - 9f_2^2)k_1^2 - 120\sigma^2 f_2 k_1 - 32\sigma^4 = 0. \tag{3.11}$$

Here, applying e_1 to both sides of (3.11), we get

$$-18k^4 + 37f_2k_1^3 + (-19f_2^2 + 530\sigma^2)k_1^2 - 440\sigma^2 f_2 k_1 - 368\sigma^4 = 0. \tag{3.12}$$

By (3.11) and (3.12), we get

$$f_2k^3 + (-202\sigma^2 + f_2^2)k_1^2 + 200\sigma^2 f_2 k_1 + 304\sigma^4 = 0. \tag{3.13}$$

Applying e_1 to both sides of (3.13) again, we get

$$f_2k_1^3 + (-f_2^2 - 604\sigma^2)k_1^2 + 606\sigma^2 f_2 k_1 + 1616\sigma^4 = 0. \tag{3.14}$$

By (3.13) and (3.14), we get

$$(201\sigma^2 + f_2^2)k_1^2 - 203\sigma^2 f_2 k_1 - 656\sigma^4 = 0. \tag{3.15}$$

By the same argument, we get

$$-f_2^2 k_1^2 + 2\sigma^2 f_2 k_1 - 354\sigma^4 = 0. \tag{3.16}$$

Substituting the 7th equation of (3.4) into (3.16), we get

$$-353\sigma^4 = c^2. \tag{3.17}$$

Applying e_1 to both sides of (3.17), we get

$$\sigma^4 k_1 = 0.$$

From this, we get

$$\sigma = 0 \text{ on } \Omega.$$

Substituting $\sigma = 0$ into (3.15), we get

$$k_1 f_2 = 0.$$

From the above arguments, we obtain $f_2 = 0$ on Ω. Then, by using the 7th equation in (3.4), we get $c = 0$ on Ω. Thus, we get $e_1(k_1) = k_1^2$. Hence, the triharmonic equation is given as

$$\triangle^M k_1^3 = 0. \tag{3.18}$$

Thus, it is easy to see that (3.18) is given by

$$k_1^5 = 0 \text{ on } \Omega.$$

Therefore, we get $k_1 = 0$ on Ω which is a contradiction. Therefore, we complete the proof of the theorem. □

4. Appendix

By the same argument as in the last section, we will give another proof of Theorem 1.1. We obtain necessary and sufficient condition for a Riemannian submersion $\varphi : (M^3(c), g) \to (N^2, h)$ to be biharmonic as follows (cf. [22]):

Corollary 4.1. *Let $\varphi : (M^3(c), g) \to (N^2, h)$ be a Riemannian submersion from a space form of constant sectional curvature c with an adapted frame $\{e_1, e_2, e_3\}$ and the integrability date f_1, f_2, k_1, k_2 and σ with $k_2 = 0$. Then, the Riemannian submersion φ is biharmonic if and only if*

$$\begin{cases} -\Delta^M k_1 + k_1(-K^N + f_1^2 + f_2^2) = 0, \\ f_1 e_1(k_1) + e_1(k_1 f_1) + f_2 e_2(k_1) + e_2(k_1 f_2) - k_1^2 f_1 = 0. \end{cases} \tag{4.1}$$

Proof of Theorem 1.1 By the same argument as in the proof of Theorem 1.3, we obtain (3.4). By the 3rd equation in (3.4), if $k_1 = 0$ on $M^3(c)$, φ is harmonic. Therefore, we consider $k_1 \neq 0$ on some set $\Omega \subset M^3(c)$ (in this case, $f_1 = 0$ on Ω) and prove that the case cannot happen.

By the 1st equation of (3.4), we get

$$e_1(\sigma) = 2k_1\sigma. \tag{4.2}$$

By the 2nd and the 7th equations in (3.4), we get

$$e_1(k_1) = k_1^2 - k_1 f_2.$$

By applying e_1 to both sides of the 7th equation in (3.4), we get

$$e_1(f_2) = f_2^2 - k_1 f_2 + 4\sigma^2.$$

Since

$$-\Delta^M k_1 = 5k_1\sigma^2 - k_1^3 - k_1 c + f_2(k_1^2 - \sigma^2 + c), \tag{4.3}$$

the biharmonic equation is given as

$$k_1^2(3\sigma^2 - k_1^2 - 3c) = 0. \tag{4.4}$$

By the assumption $k_1 \neq 0$ on Ω, we can get

$$3\sigma^2 - k_1^2 - 3c = 0. \tag{4.5}$$

Applying e_1 to this equation, we have

$$k_1^2 = 7\sigma^2 - c, \tag{4.6}$$

since $k_1 \neq 0$ on Ω. Combining (4.5), (4.6) and (4.2), we have $k_1 = \sigma = c = 0$ on Ω, which is a contradiction. Therefore, we complete the proof of the theorem. □

References

[1] Akutagawa, K., Maeta, S..: *Biharmonic properly immersed submanifolds in Euclidean spaces*. Geom. Dedicata **164** (2013), 351-355.

[2] Baird, P., Wood, J. C.: *Bernstein Theorems for harmonic Morphisms from* $\mathbb{R}^3$ *and* S^3. Math. Ann. **280**, (1988), 579-603.

[3] Baird, P., Wood, J. C.: *Harmonic Morphisms Between Riemannian Manifolds.* London Mathematical Society Monographs (N.S.) No. 29, Oxford University Press, Oxford (2003).

[4] Chen, B. Y.: *Recent developments of biharmonic conjecture and modified biharmonic conjectures*. arXiv:math.DG/1307.0245, (2013).

[5] Chen, B. Y.: *Some open problems and conjectures on submanifolds of finite type.* Soochow J. Math. **17**(2), (1991), 169-188.

[6] Defever, F.: *Hypersurfaces of* E^4 *with harmonic mean curvature vector.* Math. Nachr. **196**, (1998), 61-69.

[7] Dimitri'c, I.: *Submanifolds of Em with harmonic mean curvature vector.* Bull. Inst. Math. Acad. Sinica **20**, (1992), 53-65.

[8] Eells, J., Lemaire, L.: *Selected topics in harmonic maps*. CBMS, **50**, Amer. Math. Soc. (1983).

[9] Eells, J., Sampson, J. H.: *Harmonic mappings of Riemannian manifolds.* Amer. J. Math. **86**, (1964), 109-160.

[10] Gudmundsson, S.: *The geometry of harmonic Morphisms*. phD thesis, University of Leeds (1992).

[11] Hasanis, T., Vlachos, T.: *Hypersurfaces in* $\mathbb{E}^4$ *with harmonic mean curvature vector field.* Math. Nachr. **172**, (1995), 145-169.

[12] Jiang, G. Y.: 2*-harmonic maps and their first and second variational formula*. Chinese Ann. Math. 7A, (1986), 388–402; Note di Matematica **28**, (2009), 209–232.

[13] Luo, Y.: *Liouville-type Theorems on complete manifolds and non-existence of bi-harmonic maps*, J. Geom. Anal. **25**, (2015), 2436–2449.

[14] Maeta, S.: *k-harmonic maps into a Riemannian manifold with constant sectional curvature*. Proc. Amer. Math. Soc. **140**, (2012), 1835-1847.

[15] Maeta, S.: *Properly immersed submanifolds in complete Riemannian manifolds*, Adv. Math. **253**, (2014), 139–151.

[16] Maeta, S., Nakauchi, N., Urakawa, H.: *Triharmonic isometric immersions into a manifold of non-positively constant curvature.* Monatsh. Math, **177**, (2015), 551-567.

[17] O'Neill, B.: *The fundamental equations of a submersion.* Mich. Math. J. **13**, (1966), 459-469.
[18] Ou, Y. L.: *Some constructions of biharmonic maps and Chen's conjecture on biharmonic hypersurfaces.* J. Geom. Phys. **62**, (2012), 751-762.
[19] Ou, Y. L., G, Walschap.: *A classification of horizontally homothetic submersions from space forms of nonnegative curvature*. Bull. London Math. Soc. **38**(3), (2006), 485-493.
[20] Sakai, A.: *Riemannian Geometry*. in Japanese, Shokabo, Tokyo, (1992).
[21] Wang, S, B.: *The First Variation Formula For K-harmonic mapping*. Journal of Nanchang University **13** No 1, (1989).
[22] Wang, Z. P., Ou, Y. L.: *Biharmonic Riemannian submersions from 3-manifolds*. Math. Z. **269**, (2011) 917-925.

Department of Mathematics, Shimane University, Matsue, 690-8504, Japan.
E-mail address: tomo.miura3@gmail.com

Department of Mathematics, Shimane University, Matsue, 690-8504, Japan.
E-mail address: shun.maeta@gmail.com and maeta@riko.shimane-u.ac.jp